


TETRA-S

OVERVIEW

Tetra-S is a pure glycerine solution of the potassium salts of tetrahydro iso-alpha acids derived from CO₂ hop extract. The formulation in glycerine minimizes precipitation when exposed to cold temperatures. Even freezing during transport will not damage the product.

Tetra-S greatly enhances beer foam when used as a post-fermentation replacement for a portion of conventional bittering products.

Tetra-S provides protection against lightstruck flavor when used as the sole source for bitterness or in combination with other light stable hop products.



SPECIFICATIONS

Short description	light-stable hop extract dissolved in glycerine with intense bitterness and foam-stabilizing properties
Alpha acids	below detection limit
Iso-alpha acids	below detection limit
Tetrahydro-Iso-alpha acids	9.0 ± 0.5 % (w/w) HPLC 10.0 ± 0.5 % (w/w) Spectro
pH	9.5 ± 1.0
Density	ca. 1.22 g / ml (20°C)
Viscosity	< 3000 mPas (20° C / 68° F)

PROPERTIES

APPEARANCE

Tetra-S is a homogeneous, amber-colored, clear solution that is liquid at room temperature.

FLAVOR

Tetra-S provides 1.0 to 1.3 times the sensory bitterness compared to that achieved with iso-alpha acids. The actual intensity of bitterness depends primarily on the quantity of bittering units and the type of beer. Therefore, the target number of bittering units must be determined in preliminary tests in order to achieve the correct level of sensory bitterness.

UTILIZATION

Based on HPLC analysis of the finished beer, utilization of Tetra-S is 60 - 80 %, depending on the timing and efficiency of the addition. Actual utilization will vary from brewery to brewery due to differences in equipment and process conditions.

LIGHT STABILITY

Tetra-S only provides protection against lightstruck flavor in the complete absence of alpha acids and iso-alpha acids. Tetra-S can be used in conjunction with any Hopsteiner® light stable product to achieve light stability.

FOAM STABILITY

Tetra-S enhances both beer foam retention and cling. An improvement in beer foam is already noticeable at 2 - 3 mg/l of tetrahydro iso-alpha acids in the finished beer.

QUALITY

All Hopsteiner® products are processed in facilities which fulfill internationally recognized quality standards. A monitoring system for residues is in place.

PACKAGING

Our products are delivered in their respective recommended standard packaging. Alternatives may be possible upon customer request.

Standard packages of our processing plants in the USA (US) and Germany (DE) are:

- Canister 20 kg (US / DE)
- Jug 10 kg (US)
- Pail 20 kg (US)

USAGE

Tetra-S is typically used as a post-fermentation addition and prior to the final step in filtration.

FOR LIGHT STABLE BEER

For maximum protection against lightstruck flavor, it is essential that no other sources of non-reduced iso-alpha acids are inadvertently introduced into the wort or beer. Therefore, the following must be carefully implemented:

- exclusive use of light stable hop products throughout the entire process.
- avoid contamination through equipment surfaces previously in contact with regular iso-alpha acids.
- never pitch wort with yeast that has been in contact with regular alpha and iso-alpha acids.

DOSAGE

Dosage is based on the product concentration, an estimated or known utilization and the desired intensity of bitterness in the beer. The fact that the perceived bitterness of Tetra-S is about 1.0 to 1.3 times the bitterness of iso-alpha acids derived from conventional hop products must be taken into consideration. The correct dosage of Tetra-S must be determined through trials at the brewery.

APPLICATION

We recommend adding Tetra-S at full strength (undiluted) into the center of the beer stream for at least 70 % of the total volume being transferred. An accurate, high pressure dosing pump is required to add the product into the beer stream at a point where vigorous mixing is assured. Tetra-S can be added at ambient temperatures. If dilution is necessary, always add Tetra-S to demineralized water first and adjust the pH to 8.5 - 9.5 using either potassium hydroxide (KOH) or potassium carbonate (K₂CO₃). Laboratory scale testing is recommended prior to commercial use. If containers are used over several days, it is recommended that the headspace be flushed with nitrogen (CO₂ is not suitable).

CLEANING RECOMMENDATION

Tetra-S should not be left in dosing lines at low temperatures. Lines and dosing pumps should be flushed with warm, slightly alkaline, demineralized water or ethanol for purposes of cleaning.

STORAGE

The recommended storage temperature in the original unopened packaging is 5 - 25 °C (41 - 77 °F).

For prolonged storage, a temperature of 10 - 20 °C (50 - 68 °F) is ideal.

BEST BEFORE DATE

Under the recommended storage conditions, the shelf life from the date of production/ packaging is at least 2 years.

SAFETY

Ensure good ventilation of the workplace and wear personal protective equipment. Avoid contact with eyes and skin. Do not inhale vapors or dusts. For full safety information, please refer to the relevant Hopsteiner® safety data sheet.

ANALYTICAL METHODS

International approved methods listed in committees such as ASBC or Analytica-EBC using current standards are applied.

PRODUCT ANALYTICS

Concentration of bitter substances

- Analytica-EBC 7.9 (HPLC)
- ASBC Hops-18 (Spectro)

BEER ANALYTICS

Concentration of reduced iso-alpha acids in beer

- Analytica-EBC 9.47 (HPLC)

The standard formula for calculating bitter units in beer (Analytica-EBC 9.8 or ASBC Beer-23A) may need to be adjusted as it results in too low values when using higher amounts of Tetra-S.

Foam stability and Cling test

- NIBEM Cling
- NIBEM-T Meter
- Pour Test
- Ross & Clark
- Steinfurth Foam Stability Tester

TECHNICAL SUPPORT

We are pleased to offer assistance and advice on:

- safety data sheets
- support for brewing trials on a pilot or commercial scale
- analytical services and information about analytical procedures

Disclaimer: The information provided in this document is believed to be correct and valid. However, Hopsteiner® does not guarantee that the information provided here is complete or accurate and thus assumes no liability for any consequences resulting from its application.

Last updated: 06.05.2024

Simon H. Steiner, Hopfen, GmbH
Mainburg, Germany
Email: info(at)hopsteiner.de
Phone: +49 (0) 8751 8605 0

S.S. Steiner Inc.
New York, USA
Email: sales(at)hopsteiner.com
Phone: +(1) 212 838 8900

Steiner Hops Ltd.
Guildford, England
Email: enquiries(at)hopsteiner.co.uk
Phone: +(44) 1992 572331

Hopsteiner Trading (Zhuhai) Co., Ltd.
Zhuhai, PRC
Email: info(at)hopsteiner.com.cn
Phone: +(86) 756 322 3340